

## RENCANA PELAKSANAAN PEMBELAJARAN

SATUAN PEMBELAJARAN	: SMP NEGERI 1 Sangasanga
Mata Pelajaran	: Bahasa Inggris
Kelas / Semester	: IX / Ganjil
Materi Pokok	: Drug Label
Alokasi Waktu	: 10 Menit

### A. Tujuan Pembelajaran

- Peserta didik dapat mengidentifikasi fungsi sosial, struktur teks, dan unsur kebahasaan dari teks *label* dengan meminta dan memberi informasi terkait obat sesuai dengan konteks penggunaannya
- Peserta didik dapat menemukan informasi tersirat dan tersurat dari teks *Drug Label*, dan membuat *mindmap* atau *outline* dari teks *Drug Label*.

### B. Langkah-langkah Kegiatan Pembelajaran

Alat : Gambar Drug's label & LKPD

Bahan : Internet (<https://www.drugwatch.com/health/how-to-read-a-drug-label/>)

#### 1. Kegiatan Pembuka

- Guru membuka pembelajaran dengan salam
- Guru menanyakan kabar peserta didik dan memastikan kehadiran peserta didik
- Guru memimpin doa sebelum memulai pembelajaran
- Guru menanyakan kembali tentang pembelajaran sebelumnya
- Guru menyampaikan tujuan pembelajaran pada pertemuan tersebut

#### 2. Kegiatan Inti

- Peserta didik bekerja berpasangan untuk memasangkan kartu frasa dan kartu definisi (*Pre-Teach Vocabulary*)
- Peserta didik membaca teks dan meletakkan frase / *Drug Facts Phrase* kedalam teks rumpang dengan benar. (LKPD 1)
- Peserta didik menjawab beberapa pertanyaan dari guru (lisan) terkait fungsi sosial drug label. (Guru memancing/*eliciting* jawaban dari peserta didik)
- Peserta didik membaca dan menjawab pertanyaan-pertanyaan pada LKPD 2 secara individu.
- Peserta didik mengecek jawabannya dengan teman disampingnya (*Peer Check*).
- Guru mengecek jawaban peserta didik secara bersama-sama dan memberikan umpan balik (*whole class feedback, nominate, pounce & bounce*)

#### 3. Kegiatan Penutup

- Peserta didik menyimpulkan pembelajaran terkait dengan drug label
- Peserta didik mengungkapkan perasaannya setelah mengikuti pembelajaran hari ini
- Guru menyampaikan rencana pembelajaran berikutnya
- Guru menutup pembelajaran dengan salam

### C. Penilaian Pembelajaran

- Guru menilai LKPD
- Guru merekap penilaian sikap peserta didik selama mengikuti pembelajaran

Mengetahui,  
Kepala Sekolah

Sangasanga, 6 Januari 2022

Guru Mata Pelajaran

**Rebo, S.Pd**  
**NIP 196406151988031016**

**Rezha Hardi Pratama, S.Pd**  
NIP. -

**Pre-teaching Vocabulary**

Work in pair. Match the phrase and the definition correctly. In 5 minutes! (cut-phrases)

Phrase	Definition
<b>Active Ingredient</b>	<b>Bahan aktif penyusun obat yang membuat obat bekerja efektif</b>
<b>Uses</b>	<b>Gejala-gejala penyakit yang diobati</b>
<b>Warnings</b>	<b>Peringatan, Efek samping</b>
<b>Directions</b>	<b>Jumlah (dosis) obat, seberapa sering dikonsumsi, dan seberapa banyak obat tersebut dapat dikonsumsi secara aman dalam sehari.</b>
<b>Other Information</b>	<b>Hal-hal penting untuk diketahui seperti bagaimana cara penyimpanan obat tersebut.</b>
<b>Inactive Ingredients</b>	<b>Rasa obat, bahan pengawet, atau bahan lain yang tidak berfungsi untuk mengobati gejala penyakit.</b>
<b>Questions / Comments</b>	<b>Nomor telepon perusahaan yang dapat dihubungi jika memiliki pertanyaan / komentar tentang obat tersebut.</b>

## LKPD 1

In pairs, read the drug label and fill in the gaps with suitable phrase given.

1	<b>Purpose</b> <b>(in each caplet)</b> Acetaminophen 500 mg.....Pain reliever/fever reducer		<b>Drug Facts (continued)</b> <b>Stop use and ask a doctor if</b> <ul style="list-style-type: none"> <li>■ pain gets worse or lasts more than 10 days</li> <li>■ fever gets worse or lasts more than 3 days</li> <li>■ new symptoms occur    ■ redness or swelling is present</li> </ul> <p>These could be signs of a serious condition.</p>					
2	<ul style="list-style-type: none"> <li>■ temporarily relieves minor aches and pains due to:</li> <li>■ the common cold    ■ headache</li> <li>■ minor pain of arthritis    ■ backache</li> <li>■ muscular aches    ■ toothache</li> <li>■ premenstrual and menstrual cramps</li> <li>■ temporarily reduces fever</li> </ul>		<b>If pregnant or breast-feeding</b> , ask a health professional before use. <b>Keep out of reach of children. Overdose warning:</b> In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.					
3	<b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take <ul style="list-style-type: none"> <li>■ more than 4,000 mg of acetaminophen in 24 hours</li> <li>■ with other drugs containing acetaminophen</li> <li>■ 3 or more alcoholic drinks every day while using this product</li> </ul> <b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> <li>■ skin reddening    ■ blisters    ■ rash</li> </ul> <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use</b>    ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  <ul style="list-style-type: none"> <li>■ if you have ever had an allergic reaction to this product or any of its ingredients</li> </ul> <p><b>Ask a doctor before use if you have liver disease</b></p> <p><b>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin</b></p> </p>		<b>■ do not take more than directed (see overdose warning)</b> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="padding: 5px;">adults and children 12 years and over</td> <td style="padding: 5px;"> <ul style="list-style-type: none"> <li>■ take 2 caplets every 6 hours while symptoms last</li> <li>■ do not take more than 6 caplets in 24 hours, unless directed by a doctor</li> <li>■ do not use for more than 10 days unless directed by a doctor</li> </ul> </td> </tr> <tr> <td style="padding: 5px;">children under 12 years</td> <td style="padding: 5px;">ask a doctor</td> </tr> </table> <p>■ store at 20-25°C (68-77°F)</p>		adults and children 12 years and over	<ul style="list-style-type: none"> <li>■ take 2 caplets every 6 hours while symptoms last</li> <li>■ do not take more than 6 caplets in 24 hours, unless directed by a doctor</li> <li>■ do not use for more than 10 days unless directed by a doctor</li> </ul>	children under 12 years	ask a doctor
adults and children 12 years and over	<ul style="list-style-type: none"> <li>■ take 2 caplets every 6 hours while symptoms last</li> <li>■ do not take more than 6 caplets in 24 hours, unless directed by a doctor</li> <li>■ do not use for more than 10 days unless directed by a doctor</li> </ul>							
children under 12 years	ask a doctor							
4					5			
6					7			
Act								

Source: <https://fda.report/DailyMed/c9d3cee1-531c-439f-ae19-07acd366aad6>

Active Ingredients	Inactive ingredients	Other Information	Warnings	Uses	Questions / Comments	Directions
--------------------	----------------------	-------------------	----------	------	----------------------	------------

## LKPD 2

Individually, read the label and the statements carefully.  
Write **T** if you think the statement is true and **F** if it is false.

<b>1</b>	<b>Drug Facts</b>	<b>Drug Facts (continued)</b>				
	<b>Active ingredient (in each caplet)</b> Acetaminophen 500 mg.....Pain reliever/fever reducer	<b>Purpose</b> <b>Stop use and ask a doctor if</b> ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur   ■ redness or swelling is present These could be signs of a serious condition.				
<b>2</b>	<b>Uses</b>	<b>If pregnant or breast-feeding,</b> ask a health professional before use. <b>Keep out of reach of children. Overdose warning:</b> In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.				
<b>3</b>	<b>Warnings</b>	<b>Directions</b> <b>■ do not take more than directed (see overdose warning)</b> <table border="1"> <tr> <td>adults and children 12 years and over</td> <td>■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor</td> </tr> <tr> <td>children under 12 years</td> <td>ask a doctor</td> </tr> </table>	adults and children 12 years and over	■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor	children under 12 years	ask a doctor
adults and children 12 years and over	■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor					
children under 12 years	ask a doctor					
	<b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product  <b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening   ■ blisters   ■ rash If a skin reaction occurs, stop use and seek medical help right away.  <b>Do not use</b> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you have ever had an allergic reaction to this product or any of its ingredients  <b>Ask a doctor before use if you have liver disease</b>  <b>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin</b>	<b>Other information</b> ■ store at 20-25°C (68-77°F) <b>Inactive ingredients</b> carnauba wax, corn starch*, croscarmellose sodium*, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid *may contain one or more of these ingredients <b>Questions or comments?</b> 1-800-719-9260				

Act

Source: <https://fda.report/DailyMed/c9d3cee1-531c-439f-ae19-07acd366aad6>

No.	Statement	T / F
1	The product contains more than 4000 mg acetaminophen.	
2	Acetaminophen may cause severe skin reactions.	
3	We must store the product between 68 – 77 degree of Celcius	
4	12-year-old child must ask a doctor before consuming the product.	
5	We must stop using the product if the pain and fever get worse.	
6	We can consume it with any other drug containing acetaminophen	
7	Consuming warfarin with this medicine may be dangerous to some people	
8	Adults can consume the medicine without ask a doctor first	
9	Sodium starch glycolate in it have a medical purpose	
10	Call 1-900-719-9250 if overdose is happened	