



**PERANGKAT PEMEBALAJARAN KURIKULUM 2013
(Lesson Plan)**

**CLASS : IXa
BAHASA INGGRIS**

**OLEH
ENRIMON PIISI, S.Pd
NIM.1212321092**

**PENDIDIKAN PROFESI GURU DALAM JABATAN
PENDIDIKAN GURU BAHASA INGGRIS SMP
UNIVERSITAS KATOLIK WIDYA MANDALA SURABAYA
2021**

LESSON PLAN

School : SMPN 1 Banggai Tengah
Subject : Bahasa Inggris
Class/Semester : IX/I
Material : Medicine Label
Academic Year : 2021/2022
Time Allocation : 2 x 30 minute

A. MAIN COMPETENCE

KI 1: Concerning and practicing the value of the religion

KI 2: Exploring the attitude(honest, discipline, responsibility, care, cooperation, piece, responsive and active)and shows the attitude as problem solver in the country in effective interaction in social society and take position ourselves as a reflection of the nation in the association world.

KI 3: Understand, apply, analyze, and evaluate factual knowledge, conceptual, science, technology, art, culture, and humanities with human insight, national, state, and civilization related phenomena and events, as well as applying procedural knowledge in a specific field of study according to their talents and interests to solve the problem.

KI 4: Try, process, and serve in the concrete real and abstract domains associated with the development of the independently learned in school and home, and was able to use the method according to the rules of science.

B. Basic Competence and Indicators

Basic Competence	Indicators
3.3. Membandingkan fungsi sosial, struktur teks, dan unsur kebahasaan beberapa teks khusus dalam bentuk label, dengan meminta dan memberi informasi terkait obat/makanan/ minuman, sesuai dengan konteks penggunaannya	3.3.1 Students can identify the structure of the text on simple drug labels accurately. 3.3.2 Students can identify linguistic elements in the form of technical terms, imperatives and prohibitions from simple drug labels in an acceptable manner. 3.3.3 Students can state the social function of drug labels in an acceptable manner.
4.3 Menangkap makna secara kontekstual terkait dengan fungsi sosial, struktur teks, dan unsur kebahasaan teks khusus dalam bentuk label pendek dan sederhana, terkait obat/makanan/ minuman	4.3.1 Students can arrange random text structures about drug labels accurately. 4.3.2 Students are skilled in finding information in simple drug labels accurately.

C. LEARNING OBJECTIVE

1. Student can choose yes or no by give a checklist (√) in “Yes” column if the Label structure is available and in “No” if it is not. (C5)
2. Students can make an imperative and a prohibition based on the pictures (C6)
3. Student Can recommend the appropriate medicine for friend (C5)


D. LEARNING MATERIAL

- Variant of drugs/medicine

Pill (tablet)	Syrup/Liquid	Capsule
Ointment	Powder	

- Drug facts/Medicine label

It contains detail information about the medicine. Example:



Example:

Drug Facts									
Active ingredients	Purpose								
Medicine 100 mg.....	cough suppressant								
Medicine 150 mg.....	nasal decongestant								
Uses Temporarily relieves: ■ coughing due to minor throat and bronchial irritation ■ nasal congestion									
Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.									
When using this product ■ you may get drowsy ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children									
Stop use and seek medical help right away if allergic reaction occurs.									
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).									
Directions ■ Tablet melts in mouth. Can be taken with or without water.									
<table border="1"><thead><tr><th>Age</th><th>Dose</th></tr></thead><tbody><tr><td>adults and children 12 years and older</td><td>2 tablets every 12 hours; do not use more than 4 tablets in a 24-hour period</td></tr><tr><td>children 6 years to under 12 years</td><td>1 tablet every 12 hours; do not use more than 2 tablets in a 24-hour period</td></tr><tr><td>children under 6 years of age</td><td>ask a doctor</td></tr></tbody></table>	Age	Dose	adults and children 12 years and older	2 tablets every 12 hours; do not use more than 4 tablets in a 24-hour period	children 6 years to under 12 years	1 tablet every 12 hours; do not use more than 2 tablets in a 24-hour period	children under 6 years of age	ask a doctor	
Age	Dose								
adults and children 12 years and older	2 tablets every 12 hours; do not use more than 4 tablets in a 24-hour period								
children 6 years to under 12 years	1 tablet every 12 hours; do not use more than 2 tablets in a 24-hour period								
children under 6 years of age	ask a doctor								
Other information ■ store at 20°–25°C (68°–77°F) ■ keep dry									
Inactive ingredients anhydrous citric acid, aspartame, magnesium stearate, maltodextrin, modified food starch, sodium bicarbonate, D&C yellow no. 10									
Questions or comments? Call weekdays from 9 a.m. to 5 p.m. EST at 1-800-XXX-XXXX.									

Gambar 1



Answer these questions correctly:

1. What is/are the medicine made of?
2. What disease (s) does the medicine treat?
3. How do you consume the medicine?
4. What should you pay attention to when you consume the medicine?

- Text structure

Every medicine label/drug facts usually consists of following structure:

1. Name of the medicine/drugs
2. Active Ingredients
3. Uses
4. Warnings
5. Directions
6. Other Information
 - a. Amount/content
 - b. How to store the drug
 - c. Expiration date
7. Inactive Ingredients (colour, flavour, etc)
8. Question or comments (customer service)



Gambar 2

Example:

Drug Facts	
Active ingredients	Purpose
Medicine 100 mg.....	cough suppressant
Medicine 150 mg.....	nasal decongestant
Uses Temporarily relieves:	
<ul style="list-style-type: none"> ■ coughing due to minor throat and bronchial irritation ■ nasal congestion 	
Warnings	
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.	
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.	
When using this product	
<ul style="list-style-type: none"> ■ you may get drowsy ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children 	
Stop use and seek medical help right away if allergic reaction occurs.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).	
Directions	
■ Tablet melts in mouth. Can be taken with or without water.	
Age	Dose
adults and children 12 years and older	2 tablets every 12 hours; do not use more than 4 tablets in a 24-hour period
children 6 years to under 12 years	1 tablet every 12 hours; do not use more than 2 tablets in a 24-hour period
children under 6 years of age	ask a doctor
Other information	
<ul style="list-style-type: none"> ■ store at 20°–25°C (68°–77°F) ■ keep dry 	
Inactive ingredients	
anhydrous citric acid, aspartame, magnesium stearate, maltodextrin, modified food starch, sodium bicarbonate, D&C yellow no.10	
Questions or comments?	
Call weekdays from 9 a.m. to 5 p.m. EST at 1-800-XXX-XXXX.	

Active ingredients

Uses

Warning

Direction

Other information : how to store

Inactive ingredients

Questions or comments



Exercise 1

Read the drug facts carefully. Give a checklist (√) in “Yes” column if the structure is available and in “No” if it is not.

No	Facts	Available?	
		Yes	No
1.	Name of the drug		
2.	Active ingredients		
3.	Use (s)		
4.	Inactive ingredients		
5.	Warning		
6.	Other information:		
	a. How to store		
	b. Content/Amount		
	c. Expiration date		

7. Question/Comment
(Customer service)

Drug Facts TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL ON THE BOTTLE IS BROKEN OR MISSING.

Active ingredient (in each 5 mL teaspoonful)	Purpose
Diphenhydramine HCl 12.5 mg	Antihistamine

Uses • temporarily relieves: • runny nose • sneezing
 • itchy, watery eyes due to hay fever or other upper respiratory allergies
 • itching of the nose or throat

Warnings
Do not use • to make a child sleepy
 • if you are on a sodium-restricted diet
 • with any other product containing diphenhydramine, including one applied topically.

Ask a doctor before use if you have
 • glaucoma • trouble urinating due to an enlarged prostate gland
 • a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product • marked drowsiness may occur
 • sedatives and tranquilizers may increase drowsiness
 • excitability may occur, especially in children

Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately. ▶

Drug Facts (continued)

Directions • use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.
 • take every 4 to 6 hours
 • do not exceed 6 doses in a 24-hour period

age	dose
children 6 years to under 12 years	1 to 2 teaspoonfuls (12.5 mg to 25 mg)
children 4 years to under 6 years	do not use unless directed by a doctor
children under 4 years	do not use

Other information • each teaspoon contains: sodium 6 mg
 • store at controlled room temperature

Inactive ingredients citric acid, flavors, glycerin, poloxamer 407, purified water, red 33, red 40, sodium benzoate, sodium chloride, sodium citrate, and sugar



PROUDLY DISTRIBUTED BY:
VALU MERCHANDISERS, CO.
 KANSAS CITY, MO 64111
 Questions or Comments?
 1-866-534-4631

31-20122C

- Language features
 Example of a medicine label/drug facts

Drug Facts	
Active ingredients	Purpose
Medicine 100 mg.....	cough suppressant
Medicine 150 mg.....	nasal decongestant
Uses Temporarily relieves:	
<ul style="list-style-type: none"> ■ coughing due to minor throat and bronchial irritation ■ nasal congestion 	
Warnings	
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.	
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.	
When using this product	
<ul style="list-style-type: none"> ■ you may get drowsy ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children 	
Stop use and seek medical help right away if allergic reaction occurs.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).	
Directions	
■ Tablet melts in mouth. Can be taken with or without water.	
Age	Dose
adults and children 12 years and older	2 tablets every 12 hours; do not use more than 4 tablets in a 24-hour period
children 6 years to under 12 years	1 tablet every 12 hours; do not use more than 2 tablets in a 24-hour period
children under 6 years of age	ask a doctor
Other information	
<ul style="list-style-type: none"> ■ store at 20°–25°C (68°–77°F) ■ keep dry 	
Inactive ingredients	
anhydrous citric acid, aspartame, magnesium stearate, maltodextrin, modified food starch, sodium bicarbonate, D&C yellow no.10	
Questions or comments?	
Call weekdays from 9 a.m. to 5 p.m. EST at 1-800-XXX-XXXX.	



1. Technical terms


Vocabulary	Meaning
Drug facts	Informasi kimia
Pharmacy	Apotek
Active ingredient	Bahan aktif
Inactive ingredient	Bahan tak aktif
Warning	Peringatan
Direction	Aturan pakai
Other information	Informasi lain
Expiration date	Tanggal kadaluarsa
How to store	Cara penyimpanan
Question or comments	Pertanyaan dan komentar
Customer service	Layanan konsumen
Allergic reaction	Alergi
Disease	Penyakit
Relieve	Mengobati
Drowsy	Mengantuk
Overdose	Terlalu banyak
Nasal congestion	Hidung tersumbat

2. Imperative and prohibition

- Imperative

Formula : Infinitive + Object (+ complement)

Example :




<u>Consume</u>	<u>two tablets</u>	<u>after eating.</u>
Infinitive	Object	Complement (adverb)
<u>Get</u>	<u>medical help.</u>	
Infinitive	Object	

➤ Prohibition :

Formula : Do not/Stop + V1/Gerund(+ O +Complement)

Example :




<u>Do not</u>	<u>consume</u>	<u>after expiration date.</u>
Do not	Verb 1	Complement (adverb)
<u>Stop</u>	<u>using</u>	<u>it.</u>
Stop	Gerund	O



Exercise 2

Write an imperative and a prohibition based on the pictures.

Picture	Sentence
	Prohibition : Imperative :



• Social function of drugs labels

1. To choose the safe/appropriate product.
2. To avoid harmful effects.
3. To get the best result.
4. To give information about the product.



Exercise 3

Bayu and Sandra are sick, while Mrs. Susi is looking for a certain medicine. Can you recommend the appropriate medicine for them? Read the drug facts carefully.

I have a bronchial irritation. It makes me cough a lot. What medicine should I take?



Answer:

My eyes are itchy and watery. It seems that I have breathed something that triggers my allergy. What medicine should I take?



Answer:

Sandra

I want to improve my daughter's immune system. What medicine should I take?



Answer:

Mrs. Susi and Dhina

Compare to Vicks® DayQuil® Cough active ingredient*

HyVee

Day Time Cough Citrus Blend Flavor

All Day Cough Relief Up to 8 Hours

Dextromethorphan HBr Cough

NON-DROWSY ALCOHOL FREE ANTIHISTAMINE FREE

Cough Suppressant **12 FL OZ (355 ml)**

Do not use if printed neckband is broken or missing

Drug Facts

Active ingredient (in each 15 ml tablespoon)	Purpose
Dextromethorphan HBr 15 mg	Cough suppressant

Uses temporarily relieves cough due to minor throat and bronchial irritation

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

When using this product do not use more than directed

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

47340 2H F1

SEE NEW DOSING INFORMATION

Best Choice

Children's Allergy

Antihistamine Liquid Medication

Relieves:

- Sneezing • Runny Nose
- Itchy, Watery Eyes • Itchy Throat

Cherry Flavored

Alcohol Free Antihistamine Diphenhydramine HCl

4 FL OZ (118 mL)

Drug Facts TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL ON THE BOTTLE IS BROKEN OR MISSING.

Active ingredient (in each 5 mL teaspoonful)	Purpose
Diphenhydramine HCl 12.5 mg	Antihistamine

Uses temporarily relieves:

- runny nose
- sneezing
- itchy, watery eyes due to hay fever or other upper respiratory allergies
- itching of the nose or throat

Warnings
Do not use

- to make a child sleepy
- if you are on a sodium-restricted diet
- with any other product containing diphenhydramine, including one applied topically.

Ask a doctor before use if you have

- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness
- excitability may occur, especially in children

Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Kroger
QUALITY GUARANTEED

COMPARE TO active ingredients of
VICKS® NYQUIL® COUGH
*See back panel

NDC 30142-668-38
ALCOHOL 10%

SEE NEW DOSING DIRECTIONS

NiteTime Cough

Antihistamine • Doxylamine Succinate
Cough Suppressant • Dextromethorphan HBr

- ▶ Cough/Sneezing
- ▶ Runny Nose
- ▶ All Night Cough Relief
- ▶ Cherry Flavor

Our Pharmacists Recommend

10 FL OZ (295 mL)

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

Drug Facts

Active Ingredients (in each 15 mL tablespoon)	Purpose
Dextromethorphan HBr 15 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine

Uses temporarily relieves cold symptoms: cough
 runny nose and sneezing

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. to make a child sleepy

Ask a doctor before use if you have glaucoma a breathing problem such as emphysema or chronic bronchitis cough that occurs with too much phlegm (mucus) trouble urinating due to an enlarged prostate gland persistent or chronic cough as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product do not use more than directed
 excitability may occur, especially in children marked drowsiness may occur avoid alcoholic drinks be careful when driving a motor vehicle or operating machinery alcohol, sedatives and tranquilizers may increase drowsiness

SEE NEW WARNING & DOSING INFORMATION

CVS
pharmacy

NDC 49580-2464-4

NEW!

Dye-free
Children's
COLD RELIEF
SUSPENSION LIQUID

PAIN RELIEVER/FEVER REDUCER/ANTIHISTAMINE/
NASAL DECONGESTANT
Acetaminophen/Chlorpheniramine Maleate/
Phenylephrine HCl

Grape Flavor

Non-Staining Dye-Free

For ages 4 to 11
4 FL OZ (120 mL)

Drug Facts

Active ingredients (in each 5 mL, 1 teaspoon)	Purpose
Acetaminophen 160 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 1 mg	Antihistamine
Phenylephrine HCl 2.5 mg	Nasal decongestant

Uses temporarily relieves the following cold/flu symptoms: • minor aches and pains • headache • sore throat • sneezing and runny nose • stuffy nose • temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: • more than 5 doses in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • to make a child sleepy • in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product. • with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if the child has • liver disease • glaucoma • heart disease • diabetes • high blood pressure • thyroid disease • a breathing problem such as chronic bronchitis

Ask a doctor or pharmacist before use if the child is taking • sedatives or tranquilizers • the blood thinning drug warfarin

When using this product • do not exceed recommended dosage (see overdose warning) • may cause excitability, especially in children • marked drowsiness may occur • sedatives and tranquilizers may increase drowsiness

SEE NEW DOSING INFORMATION

HEALTH CARE
Best Choice Brand

Children's Non-Aspirin

Acetaminophen Oral Suspension

Fever Reducer • Pain Reliever

- Alcohol Free
- Aspirin Free
- Ibuprofen Free

GRAPE FLAVOR

4 FL OZ (118 mL)

Drug Facts

Active ingredient (in each 5 mL teaspoonful)	Purpose
Acetaminophen 160 mg	Pain reliever/fever reducer

Uses temporarily: • reduces fever • relieves minor aches and pains due to: • the common cold • flu • headaches • sore throat • immunizations

Warnings
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other acetaminophen containing products. This may lead to an overdose, which may cause liver damage. (see overdose warning)

When using this product • do not exceed recommended dosage (see overdose warning)

Stop use and ask a doctor if • new symptoms occur • redness or swelling is present • pain gets worse or lasts for more than 5 days • fever gets worse or lasts for more than 3 days

Keep this and all drugs out of the reach of children.
Overdose Warning: Taking more than the recommended dose (overdose) could cause serious health problems, including liver damage. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately. Quick medical attention is critical even if you do not notice any signs or symptoms.

PROUDLY DISTRIBUTED BY:
VALU MERCHANDISERS, CO.
KANSAS CITY, MO 64111
Questions or Comments?
1-866-534-4631

Best Choice
100% GUARANTEED
www.awgbrands.com



Exercise 4

Instruction:

- Make a group of 4-5 students.
- Arrange the papers you have got into a good drug facts (pay attention to the text structure).
- Draw some lines to identify each part of the text. (Look at the example)
- Read your drug facts carefully and then answer these questions:

Questions :

1. What is/are the medicine made of?
2. What disease (s) does the medicine treat?
3. How do you consume the medicine?
4. What should you pay attention to when you consume the medicine?

Drug Facts:

Drug Facts

Active ingredient (in each 15 ml tablespoon)	Purpose
Dextromethorphan HBr 15 mg.....	Cough suppressant

Uses temporarily relieves cough due to minor throat and bronchial irritation

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

When using this product do not use more than directed

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Drug Facts (continued)

Directions

- take only as recommended
- use dose cup
- do not exceed 4 doses per 24 hours

adults and children 12 years and over	30 ml (2 TBSP) every 6 to 8 hours
children 6 to under 12 years	15 ml (1 TBSP) every 6 to 8 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

when using other Daytime or Nighttime products, carefully read each label to insure correct dosing

Other information

- each tablespoon contains: sodium 13 mg
- store at 20°-25°C (68°-77°F)

Inactive ingredients anhydrous citric acid, D&C yellow #10, FD&C yellow #6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments? 1-800-719-9260

*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® DayQuil® Cough.


DISTRIBUTED BY HY-VEE, INC.
 5820 WESTOWN PARKWAY
 WEST DES MOINES, IA 50266
 SATISFACTION GUARANTEED
 OR PURCHASE PRICE REFUNDED
 FOR INFORMATION CALL
 1-800-289-8343
 www.hy-vee.com

GLUTEN FREE



- Homework

Find a simple medicine label/drug facts. Read the fact carefully and then identify each part of the text by creating a diagram below.



- Remedial Materials

Read the drug facts carefully. Give a checklist (✓) in “Yes” column if the structure is available and in “No” column if it is not.

No	Facts	Available?	
		Yes	No
1.	Name of the drug		
2.	Active ingredients		
3.	Use (s)		
4.	Inactive ingredients		
5.	Warning		
6.	Other information: a. How to store b. Content/Amount c. Expiration date		
7.	Question/Comment (Customer service)		

Drug Facts TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL ON THE BOTTLE IS BROKEN OR MISSING.

Active ingredient (in each 5 mL teaspoonful) Purpose
Diphenhydramine HCl 12.5 mgAntihistamine

Uses • temporarily relieves: • runny nose • sneezing
• itchy, watery eyes due to hay fever or other upper respiratory allergies
• itching of the nose or throat

Warnings
Do not use • to make a child sleepy
• if you are on a sodium-restricted diet
• with any other product containing diphenhydramine, including one applied topically.

Ask a doctor before use if you have
• glaucoma • trouble urinating due to an enlarged prostate gland
• a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product • marked drowsiness may occur
• sedatives and tranquilizers may increase drowsiness
• excitability may occur, especially in children

Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Drug Facts (continued)

Directions • use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.
• take every 4 to 6 hours
• do not exceed 6 doses in a 24-hour period

age	dose
children 6 years to under 12 years	1 to 2 teaspoonfuls (12.5 mg to 25 mg)
children 4 years to under 6 years	do not use unless directed by a doctor
children under 4 years	do not use

Other information • each teaspoon contains: sodium 6 mg
• store at controlled room temperature

Inactive ingredients citric acid, flavors, glycerin, poloxamer 407, purified water, red 33, red 40, sodium benzoate, sodium chloride, sodium citrate, and sugar



PROUDLY DISTRIBUTED BY:
VALU MERCHANDISERS, CO.
KANSAS CITY, MO 64111
Questions or Comments?
1-866-534-4631

31-20122C

• Enrichment Materials

Susanto and Tajudin are sick, while Dora is searching for a correct medicine. Can you give the appropriate medicine for them? Read the drug facts carefully.


My throat is so sore. It makes me cough for several days. What should I take to relief it?



Susanto

Answer:

Urgh, there is no water here. Can you please help me? I need to eat my lunch.



Answer:

Dora

My body aches and I have a fever too. What medicine should I take?



Answer:

Tajudin

SINA - SPRITZ
INSTANT HAND DEGERMER

Kills 99.99% of *E. coli*, *Salmonella enterica* and *Staphylococcus aureus* (MRSA) in 15 seconds.

ENHANCED WITH MOISTURIZERS

DANGER: FLAMMABLE
KEEP OUT OF REACH OF CHILDREN
KEEP AWAY FROM FIRE OR FLAME
FOR EXTERNAL USE ONLY

See other cautions on opposite panel of label

NET CONTENTS: 4 FL. OZ. (118 mL)

LEADER
NDC 37205-719-34
SEE NEW WARNINGS

Compare to Vicks® DayQuil® active ingredients

Non-Drowsy
Day•Time
Cold & Flu
Multi-Symptom Relief

Acetaminophen, Phenylephrine HCl, Dextromethorphan HBr
Cough Suppressant, Nasal Decongestant, Fever Reducer, Pain Reliever

ALCOHOL FREE
ANTIHISTAMINE FREE

8 FL OZ (237 mL)

Drug Facts

Active Ingredients (in each 15 mL) **Purpose**
Acetaminophen 325 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....Cough suppressant
Phenylephrine HCl 5 mg.....Nasal decongestant

Uses
temporarily relieves common cold/flu symptoms:
headache sore throat fever
cough due to minor throat and bronchial irritation
minor aches and pains nasal congestion

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if
adult takes more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
child takes more than 4 doses (15 mL each) in 24 hours, which is the maximum daily amount for this product
taken with other drugs containing acetaminophen
adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
skin reddening blisters rash
If a skin reaction occurs, stop use and seek medical help right away.

PEEL BACK AT CORNER FOR MORE INFORMATION

Drug Facts (continued)

Directions
do not exceed recommended dosage (see overdose warning)
shake well before using
find right dose on chart below. If possible, use weight to dose; otherwise, use age.
if needed, repeat dose every 4 hours
do not use more than 5 times in 24 hours
only use enclosed measuring cup

Weight (lb)	Age (yr)	Dose (tsp or mL)
under 36	under 4	do not use
36-48	4-6	do not use unless directed by a doctor
49-59	7-8	2 tsps or 10 mL
60-71	9-10	2 1/2 tsps or 12.5 mL
72-95	11	3 tsps or 15 mL

Other information • dosage cup provided
store at controlled room temperature

Inactive Ingredients
blue 1, butyl paraben, carboxymethylcellulose sodium, cellulose, citric acid, flavors, glycerin, high fructose corn syrup, propylene glycol, purified water, red 33, sodium benzoate, sorbitol, sucralose, xanthan gum

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL ON THE BOTTLE IS BROKEN OR MISSING.

31-58651C

SEE NEW DOSING INFORMATION

HEALTH CARE
Best Choice Brand

Children's Non-Aspirin
Acetaminophen Oral Suspension

Fever Reducer • Pain Reliever
• Alcohol Free
• Aspirin Free
• Ibuprofen Free

GRAPE FLAVOR

4 FL OZ (118 mL)

Drug Facts

Active ingredient (in each 5 mL teaspoonful) **Purpose**
Acetaminophen 160 mg.....Pain reliever/fever reducer

Uses temporarily: • reduces fever • relieves minor aches and pains due to:
the common cold • flu • headaches • sore throat • immunizations • toothaches

Warnings
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use • with any other acetaminophen containing products. This may lead to an overdose, which may cause liver damage. (see overdose warning)

When using this product
do not exceed recommended dosage (see overdose warning)

Stop use and ask a doctor if
new symptoms occur • redness or swelling is present
pain gets worse or lasts for more than 5 days
fever gets worse or lasts for more than 3 days

Keep this and all drugs out of the reach of children.
Overdose Warning: Taking more than the recommended dose (overdose) could cause serious health problems, including liver damage. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately. Quick medical attention is critical even if you do not notice any signs or symptoms.

PROUDLY DISTRIBUTED BY:
VALU MERCHANTISERS, CO.
KANSAS CITY, MO 64111
Questions or Comments?
1-866-534-4631

Best Choice
100% GUARANTEED
www.awgbrands.com



E. Metode Pembelajaran

- Discovery-based learning

Scientific Approach	Discovery-Based Learning
Mengamati	Perumusan masalah
Menanya	
Mengumpulan informasi	Hipotesis Pengumpulan data
Mengasosiasi	Pengumpulan data
Mengkomunikasikan	Komunikasi

E. Media Pembelajaran

- Power Point
- Handout
- Realia (bungkus obat)
- Pictures
- Karton warna

F. Sumber Belajar

- Reading Medicine Labels and Instruction. URL : <http://esol.britishcouncil.org/content/learners/uk-life/be-safe-uk/reading-medicine-labels-and-instructions> (Diunduh pada 31 Maret 2017)
- How to Read Medicine Labels. URL : <https://www.tylenol.com/safety-dosing/usage/how-to-read-medicine-labels> (Diunduh pada 03 April 2017)
- Medicine labels. URL : <https://esol.britishcouncil.org/sites/default/files/medicine%20labels.pdf> (Diunduh pada 31 Maret 2017)
- Reading Drug Facts. URL : <http://www.scholastic.com/browse/article.jsp?id=3757745> (Diunduh pada 31 Maret 2017)

- O'Brien, Elizabeth. *Imperative Sentence*. URL : <http://www.english-grammar-revolution.com/imperative-sentence.html> (Diunduh pada 26 Maret 2017)
- Modals. URL : <https://learnenglish.britishcouncil.org/en/quick-grammar/modals-1> (Diunduh pada 30 Maret 2017)
- Wachidah, Siti. Khatimah, Y.R.,Diyantari. 2015. *Bahasa Inggris, Think Globally Act Locally*. Jakarta : Kementrian Pendidikan dan Kebudayaan.
- Children's Allergy. URL : <https://www.drugs.com/pro/children-s-allergy.html> (Diunduh pada 01 April 2017)
- Cough Suppressant. URL : <https://dailymed.nlm.nih.gov/dailymed/archives/fdaDrugInfo.cfm?archiveid=55954> (Diunduh pada 01 April 2017)
- Medicine bottles. URL : http://article.images.consumerreports.org/prod/content/dam/CRE/salud/medicinas/costco_s (Diunduh pada 29 Maret 2017)

G. LEARNING ACTIVITIES

First Meeting (2 x 30 minutes)

No	ACTIVITIES	Description	Component 4c or PPK	Level HOTS	Time
1	Pre Interactional activity	The teachers says greeting to greet the students in classroom	(Integrity)		10 minutes
		Students respond the teacher's greeting			
		The class continued with pray together, one of student lead to pray before starting the lesson	(Independently, Confidence, Religious)		
		Sing Indonesia Raya anthem	(Nationalist)		
		The Teacher check students attendance	(Discipline)		
		The teacher remind the students to <ol style="list-style-type: none"> 1. Wear mask to protect yourself from covid 19 2. Clean your hand often with soap 3. Stay at last seven feet from other 			
		Students answer questions about the material that has been studied previously with the material to be studied.	(apperception)		
		Students listen to the teacher's explanation that relates the material	(apperception)		

		(Data collection)			
		1. Students compose jumbled text about drug labels in groups. 2. Students look for information on drug labels by answering questions in groups. 3. Students collect the results of the discussion in the form of jumbled text that has been compiled and answers. (Communicating)			
3	Post Teaching	Students listen to the feedback given by the teacher carefully			10 minutes
		Students conclude the material by mentioning what material they have learned that day.			
		Students record learning activities for the next meeting.			
		The teacher close the learning activity by inviting students pray before ending the lesson	Religious		
		The student respond the teacher greeting			

H. Penilaian Hasil Belajar

- Penilaian Sikap
 1. Teknik penilaian : observasi
 2. Bentuk penulisan : skoring
 3. Instrumen penilaian :

No	Nama Siswa	Aspek		
		Berani	Percaya Diri	Kerja Sama
1.				
2.				
3.				
dst				
Total				

4. Rubrik penilaian

Berani	
Kriteria	Skor
Menunjukkan sikap berani	5
Beberapa kali menunjukkan sikap berani (1-2)	4
Pernah menunjukkan sikap berani (3-4)	3
Sering menunjukkan sikap tidak berani (5-6)	2
Sangat sering menunjukkan sikap tidak berani	1

Percaya diri	
Kriteria	Skor
Menunjukkan sikap percaya diri	5
Beberapa kali menunjukkan sikap tidak percaya diri (1-2)	4
Pernah menunjukkan sikap tidak percaya diri (3-4)	3
Sering menunjukkan sikap tidak percaya diri (5-6)	2
Sangat sering menunjukkan sikap tidak percaya diri	1

Kerja Sama	
Kriteria	Skor
Selalu bekerja sama	5
Sering bekerja sama (5-6)	4
Beberapa kali melakukan kerja sama (3-4)	3
Pernah bekerja sama (1-2)	2
Tidak pernah bekerja sama	1

Nilai : (Total skor x 2) : 3

Contoh :

- Berani : 5
- Percaya diri : 5
- Kerja sama : 4
- Total skor : 14

Maka Nilai : (total skor x 2) : 3
 : (14 x 2) : 3 = 9,3

• Penilaian Pengetahuan

1. Teknik penilaian : test tertulis
2. Bentuk penulisan : identifikasi teks
 membuat kalimat
 menjodohkan gambar dengan teks
3. Kisi-kisi :

No	Kompetensi Dasar	Materi	Indikator soal	Bentuk soal	Jumlah soal
1.	3.3 Membandingkan fungsi sosial, struktur teks, dan unsur kebahasaan beberapa teks khusus dalam bentuk label, dengan meminta dan memberi informasi terkait obat/makanan/minuman, sesuai dengan konteks penggunaannya	Label pada obat	3.3.1 Siswa dapat mengidentifikasi struktur teks yang ada pada label obat sederhana secara akurat.	Mengidentifikasi kasi struktur teks	10
			3.3.2 Siswa dapat mengidentifikasi unsur kebahasaan berupa <i>technical term, imperative and prohibition</i> dari label obat sederhana secara akurat.	Membuat kalimat (<i>imperative and prohibition</i>)	4
			3.3.3 Siswa dapat menyebutkan fungsi sosial dari label obat secara berterima.	Menjodohkan gambar dengan teks	3

4. Instrumen penilaian :



Exercise 1

Read the drug facts carefully. Give a checklist (√) in “Yes” column if the structure is available and in “No” if it is not.

Drug Facts TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL ON THE BOTTLE IS BROKEN OR MISSING.

Active ingredient (in each 5 mL teaspoonful) Purpose
 Diphenhydramine HCl 12.5 mgAntihistamine

Uses • temporarily relieves: • runny nose • sneezing
 • itchy, watery eyes due to hay fever or other upper respiratory allergies
 • itching of the nose or throat

Warnings
Do not use • to make a child sleepy
 • if you are on a sodium-restricted diet
 • with any other product containing diphenhydramine, including one applied topically.

Ask a doctor before use if you have
 • glaucoma • trouble urinating due to an enlarged prostate gland
 • a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product • marked drowsiness may occur
 • sedatives and tranquilizers may increase drowsiness
 • excitability may occur, especially in children

Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Drug Facts (continued)

Directions • use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.
 • take every 4 to 6 hours
 • do not exceed 6 doses in a 24-hour period

age	dose
children 6 years to under 12 years	1 to 2 teaspoonfuls (12.5 mg to 25 mg)
children 4 years to under 6 years	do not use unless directed by a doctor
children under 4 years	do not use

Other information • each teaspoon contains: sodium 6 mg
 • store at controlled room temperature

Inactive ingredients citric acid, flavors, glycerin, poloxamer 407, purified water, red 33, red 40, sodium benzoate, sodium chloride, sodium citrate, and sugar



PROUDLY DISTRIBUTED BY:
VALU MERCHANDISERS, CO.
KANSAS CITY, MO 64111
 Questions or Comments?
 1-866-534-4631

31-20122C



No	Facts	Available?	
		Yes	No
1.	Name of the drug		
2.	Active ingredients		
3.	Use (s)		
4.	Inactive ingredients		
5.	Warning		
6.	Other information: a. How to store b. Content/Amount c. Expiration date		
7.	Question/Comment (Customer service)		



Exercise 2

Write an imperative and a prohibition based on the pictures.

Picture	Sentence
	Prohibition :
	Imperative :

Prohibition:

Imperative:



Exercise 3

Bayu and Sandra are sick, while Mrs. Susi is looking for a certain medicine. Can you recommend the appropriate medicine for them? Read the drug facts carefully.

<p>I have a bronchial irritation. It makes me cough a lot. What medicine should I take?</p>	 <p>Bayu</p>	<p>Answer:</p>
---	---	----------------

<p>My eyes are itchy and watery. It seems that I breathed something that triggers my allergy. What medicine should I take?</p>	 <p>Sandra</p>	<p>Answer:</p>
--	---	----------------

<p>I want my daughter to improve my daughter's immune system. What medicine should I take?</p>		<p>Answer:</p>
--	---	----------------

Mrs. Susi and Dhina



COMPARE TO active ingredients of
VICKS® NYQUIT® COUGH
*See back panel

NDC 30142-668-38
ALCOHOL 10%

SEE NEW DOSING DIRECTIONS

NiteTime Cough

**Antihistamine • Doxylamine Succinate
Cough Suppressant • Dextromethorphan HBr**

- ▶ Cough/Sneezing
- ▶ Runny Nose
- ▶ All Night Cough Relief
- ▶ Cherry Flavor

**Our
Pharmacists
Recommend**

10 FL OZ (295 mL)

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each 15 mL tablespoon)	Purpose
Dextromethorphan HBr 15 mg.....	Cough suppressant
Doxylamine succinate 6.25 mg.....	Antihistamine

Uses temporarily relieves cold symptoms: ■ cough
■ runny nose and sneezing

Warnings
Do not use ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ to make a child sleepy

Ask a doctor before use if you have ■ glaucoma ■ a breathing problem such as emphysema or chronic bronchitis ■ cough that occurs with too much phlegm (mucus) ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product ■ do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives and tranquilizers may increase drowsiness

SEE NEW DOSING INFORMATION



Children's Allergy

**Antihistamine
Liquid Medication**

Relieves:

- Sneezing • Runny Nose
- Itchy, Watery Eyes • Itchy Throat

Cherry Flavored

Alcohol Free
Antihistamine
Diphenhydramine HCl

4 FL OZ (118 mL)

Drug Facts **TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL ON THE BOTTLE IS BROKEN OR MISSING.**

Active ingredient (in each 5 mL teaspoonful)	Purpose
Diphenhydramine HCl 12.5 mg	Antihistamine

Uses • temporarily relieves: • runny nose • sneezing
• itchy, watery eyes due to hay fever or other upper respiratory allergies
• itching of the nose or throat

Warnings
Do not use • to make a child sleepy
• if you are on a sodium-restricted diet
• with any other product containing diphenhydramine, including one applied topically.

Ask a doctor before use if you have
• glaucoma • trouble urinating due to an enlarged prostate gland
• a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product • marked drowsiness may occur
• sedatives and tranquilizers may increase drowsiness
• excitability may occur, especially in children

Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

SEE NEW WARNING & DOSING INFORMATION

CVS
pharmacy

NDC 49580-2464-4

NEW!

Dye-free
Children's
**COLD
RELIEF**

SUSPENSION LIQUID

PAIN RELIEVER/FEVER REDUCER/ANTIHISTAMINE/
NASAL DECONGESTANT

**Acetaminophen/Chlorpheniramine Maleate/
Phenylephrine HCl**

Grape Flavor

**Non-Staining
Dye-Free**



For ages 4 to 11

4 FL OZ (120 mL)

Drug Facts

Active ingredients (in each 5 mL, 1 teaspoon)	Purpose
Acetaminophen 160 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 1 mg	Antihistamine
Phenylephrine HCl 2.5 mg	Nasal decongestant

Uses temporarily relieves the following cold/flu symptoms: • minor aches and pains • headache • sore throat • sneezing and runny nose • stuffy nose • temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: • more than 5 doses in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • to make a child sleepy • in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product. • with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if the child has • liver disease • glaucoma • heart disease • diabetes • high blood pressure • thyroid disease • a breathing problem such as chronic bronchitis

Ask a doctor or pharmacist before use if the child is taking • sedatives or tranquilizers • the blood thinning drug warfarin

When using this product • do not exceed recommended dosage (see overdose warning) • may cause excitability, especially in children • marked drowsiness may occur • sedatives and tranquilizers may increase drowsiness

NDC 42507-473-40

Compare to Vicks® DayQuil® Cough
active ingredient*

HyVee

**Day Time
Cough**

Citrus
Blend
Flavor



**All Day Cough Relief
Up to 8 Hours**

Dextromethorphan HBr Cough

**NON-DROWSY
ALCOHOL FREE
ANTIHISTAMINE FREE**

Cough Suppressant

**12 FL OZ
(355 ml)**

Do not use if printed neckband is broken or missing

Drug Facts

Active ingredient (in each 15 ml tablespoon)	Purpose
Dextromethorphan HBr 15 mg	Cough suppressant

Uses temporarily relieves cough due to minor throat and bronchial irritation

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
■ cough that occurs with too much phlegm (mucus)
■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema

When using this product do not use more than directed

Stop use and ask a doctor if
■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

47340 2H F1

SEE NEW DOSING INFORMATION

HEALTH CARE
Best Choice Brand

Children's Non-Aspirin
Acetaminophen Oral Suspension

Fever Reducer • Pain Reliever

- Alcohol Free
- Aspirin Free
- Ibuprofen Free

GRAPE FLAVOR

4 FL OZ (118 mL)

Drug Facts

Active ingredient (in each 5 mL teaspoonful)	Purpose
Acetaminophen 160 mg.....	Pain reliever/fever reducer

Uses temporarily: • reduces fever • relieves minor aches and pains due to:
• the common cold • flu • headaches • sore throat • immunizations

Warnings
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other acetaminophen containing products. This may lead to an overdose, which may cause liver damage. (see overdose warning)

When using this product
• do not exceed recommended dosage (see overdose warning)

Stop use and ask a doctor if
• new symptoms occur • redness or swelling is present
• pain gets worse or lasts for more than 5 days
• fever gets worse or lasts for more than 3 days

Keep this and all drugs out of the reach of children.
Overdose Warning: Taking more than the recommended dose (overdose) could cause serious health problems, including liver damage. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately. Quick medical attention is critical even if you do not notice any signs or symptoms.

PROUDLY DISTRIBUTED BY:
VALU MERCHANDISERS, CO.
KANSAS CITY, MO 64111
Questions or Comments?
1-866-534-4631

Best Choice
100% GUARANTEED
www.awgbrands.com

5. Rubrik penilaian :

- Exercise 1

Kriteria	Skor
Jawaban benar	10
Jawaban salah	0

Skor : jawaban benar x 100

Contoh:

Jawaban benar 10

Jumlah soal 10

Maka : $10 \times 10 = 100$

- Exercise 2

Aspek	Kriteria	Skor
Kesesuaian jawaban dengan pertanyaan	• Jawaban sesuai dengan pertanyaan dan teks yang diberikan.	4
	• Separuh jawaban tidak sesuai dengan pertanyaan dan teks yang diberikan.	2
	• Jawaban tidak sesuai dengan materi yang dibahas.	1
Tata Bahasa	• Semua kalimat disusun dengan tata bahasa yang tepat.	4
	• Beberapa kata dalam kalimat disusun dengan tata bahasa yang kurang tepat. (2-3)	2

	<ul style="list-style-type: none"> • Kalimat tidak disusun dengan tata bahasa yang tepat. 	1
Ejaan	<ul style="list-style-type: none"> • Kalimat ditulis dengan ejaan yang benar. • Beberapa kata ditulis dengan ejaan yang kurang tepat. (2-3) • Tidak ada kata yang ditulis dengan ejaan yang tepat. 	4 2 1
Tanda baca	<ul style="list-style-type: none"> • Kalimat ditulis dengan tanda baca yang tepat. • Beberapa kata ditulis dengan tanda baca yang kurang tepat. (<i>capitalization</i>) (1-2) • Tidak ada kalimat yang ditulis dengan tanda baca yang tepat. 	4 2 1

• Exercise 3

Kriteria	Skor
Jawaban benar	10
Jawaban salah	0

$$\text{Skor} : \frac{\text{jawaban benar} \times 100}{\text{Jumlah soal}}$$

Contoh:

Jawaban benar 3

Jumlah soal 3

Maka : $(3 \times 100) : 3 = 100$

6. Kunci jawaban :

Exercise 1

No	Facts	Available?	
		Yes	No
1.	Name of the drug	√	
2.	Active ingredients	√	
3.	Use (s)	√	
4.	Inactive ingredients	√	
5.	Warning	√	
6.	Other information:		
	a. How to store	√	
	b. Content/Amount	√	
	c. Expiration date		√
7.	Question/Comment (Customer service)	√	

Exercise 2

1. Prohibition : Do not drink using spoon.
Imperative : Drink the medicine using the measure cup.
2. Prohibition : Do not drink directly.

Imperative : Shake well before using.

Exercise 3

1. HyVee Day Time
2. Children's Allergy
3. Children's non-aspirin

- Penilaian Keterampilan

1. Teknik penilaian : test tertulis
2. Bentuk penilaian : pertanyaan
3. Kisi-kisi :

No	Kompetensi Dasar	Materi	Indikator soal	Bentuk soal	Jumlah soal
1.	4.3Menangkap makna secara kontekstual terkait dengan fungsi sosial, struktur teks, dan unsur kebahasaan teks khusus dalam bentuk label pendekdansederhana, terkait obat/makanan/minuman	Label pada obat	4.3.1 Siswa terampil menemukan informasi dalam label obat sederhana secara akurat.	Pertanyaan	7

4. Instrumen penilaian :



Exercise 4


Instruction:

- Make a group of 4-5 students.
- Arrange the papers you have got into a good drug facts (pay attention to the text structure).
- Draw some lines to identify each part of the text. (Look at the example)
- Present your drug facts in front of the class by answering these questions:

Questions :

1. What is/are the medicine made of?
2. What disease (s) does the medicine treat?
3. How do you consume the medicine?
4. What should you pay attention to when you consume the medicine?

Drug facts :

<p>Drug Facts</p> <p>Active ingredient (in each 15 ml tablespoon) Purpose Dextromethorphan HBr 15 mg.....Cough suppressant</p>									
<p>Uses temporarily relieves cough due to minor throat and bronchial irritation</p>									
<p>Warnings Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p>									
<p>Ask a doctor before use if you have ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema</p>									
<p>When using this product do not use more than directed</p>									
<p>Stop use and ask a doctor if ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.</p>									
<p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p>									
<p>Drug Facts (continued)</p> <p>Directions ■ take only as recommended ■ use dose cup ■ do not exceed 4 doses per 24 hours</p> <table border="1"> <tr> <td>adults and children 12 years and over</td> <td>30 ml (2 TBSP) every 6 to 8 hours</td> </tr> <tr> <td>children 6 to under 12 years</td> <td>15 ml (1 TBSP) every 6 to 8 hours</td> </tr> <tr> <td>children 4 to under 6 years</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 years</td> <td>do not use</td> </tr> </table> <p>■ when using other Daytime or Nighttime products, carefully read each label to insure correct dosing</p>		adults and children 12 years and over	30 ml (2 TBSP) every 6 to 8 hours	children 6 to under 12 years	15 ml (1 TBSP) every 6 to 8 hours	children 4 to under 6 years	ask a doctor	children under 4 years	do not use
adults and children 12 years and over	30 ml (2 TBSP) every 6 to 8 hours								
children 6 to under 12 years	15 ml (1 TBSP) every 6 to 8 hours								
children 4 to under 6 years	ask a doctor								
children under 4 years	do not use								
<p>Other information ■ each tablespoon contains: sodium 13 mg ■ store at 20°-25°C (68°-77°F)</p>									
<p>Inactive ingredients anhydrous citric acid, D&C yellow #10, FD&C yellow #6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate</p>									
<p>Questions or comments? 1-800-719-9260</p>									
<p>*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® DayQuil® Cough.</p>									
<p>DISTRIBUTED BY HY-VEE, INC. 5820 WESTOWN PARKWAY WEST DES MOINES, IA 50266 SATISFACTION GUARANTEED OR PURCHASE PRICE REFUNDED FOR INFORMATION CALL 1-800-289-8343 www.hy-vee.com</p>									
<p style="text-align: center;">GLUTEN FREE</p> 									

5. Rubrik penilaian :
Exercise 4

- Jumbled text

Aspek	Kriteria	Skor
Struktur teks	• Seluruh struktur teks disusun dengan baik sesuai dengan materi yang dibahas.	4
	• Beberapa struktur dalam teks tidak sesuai dengan struktur teks yang dibahas. (1-2)	3
	• Separuh dari seluruh stuktur teks tidak sesuai dengan struktur teks yang dibahas. (3-4)	2
	• Struktur teks tidak sesuai dengan struktur teks yang dibahas.	1

- Questions (tiap nomor)

Aspek	Kriteria	Skor
Kesesuaian jawaban dengan	• Jawaban sesuai dengan pertanyaan dan teks yang diberikan.	4

pertanyaan	• Separuh jawaban tidak sesuai dengan pertanyaan dan teks yang diberikan.	2
	• Jawaban tidak sesuai dengan materi yang dibahas.	1
Tata Bahasa	• Semua kalimat disusun dengan tata bahasa yang tepat.	4
	• Beberapa kata dalam kalimat disusun dengan tata bahasa yang kurang tepat. (2-3)	2
	• Kalimat tidak disusun dengan tata bahasa yang tepat.	1
Ejaan	• Kalimat ditulis dengan ejaan yang benar.	4
	• Beberapa kata ditulis dengan ejaan yang kurang tepat. (2-3)	2
	• Tidak ada kata yang ditulis dengan ejaan yang tepat.	1
Tanda baca	• Kalimat ditulis dengan tanda baca yang tepat.	4
	• Beberapa kata ditulis dengan tanda baca yang kurang tepat. (<i>capitalization</i>) (1-2)	2
	• Tidak ada kalimat yang ditulis dengan tanda baca yang tepat.	1


Total Nilai : Total skor x 5

Contoh :

Aspek	Nilai
Struktur text	4
Kesesuaian jawaban	4
Ejaan	4
Tanda baca	4
Tata bahasa	4
Total	20

Maka nilai : $20 \times 5 = 100$

6. Kunci jawaban :

Drug Facts		Drug Facts (continued)	
Active ingredient (in each 15 ml tablespoon)	Purpose	Directions	
Dextromethorphan HBr 15 mg.....	Cough suppressant	<ul style="list-style-type: none"> ■ take only as recommended ■ use dose cup ■ do not exceed 4 doses per 24 hours 	
Uses temporarily relieves cough due to minor throat and bronchial irritation		adults and children 12 years and over	30 ml (2 TBSP) every 6 to 8 hours
Warnings Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.		children 6 to under 12 years	15 ml (1 TBSP) every 6 to 8 hours
Ask a doctor before use if you have		children 4 to under 6 years	ask a doctor
<ul style="list-style-type: none"> ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema 		children under 4 years	do not use
When using this product do not use more than directed		<ul style="list-style-type: none"> ■ when using other Daytime or Nighttime products, carefully read each label to insure correct dosing 	
Stop use and ask a doctor if		Other information	
<ul style="list-style-type: none"> ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition. 		<ul style="list-style-type: none"> ■ each tablespoon contains: sodium 13 mg ■ store at 20°-25°C (68°-77°F) 	
If pregnant or breast-feeding, ask a health professional before use.		Inactive ingredients anhydrous citric acid, D&C yellow #10, FD&C yellow #6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)		<ul style="list-style-type: none"> ■ when using other Daytime or Nighttime products, carefully read each label to insure correct dosing 	
		Questions or comments? 1-800-719-9260	
		*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® DayQuil® Cough.	
		DISTRIBUTED BY HY-VEE, INC. 5820 WESTOWN PARKWAY WEST DES MOINES, IA 50266 SATISFACTION GUARANTEED OR PURCHASE PRICE REFUNDED FOR INFORMATION CALL 1-800-288-8343 www.hy-vee.com	
		GLUTEN FREE	
			

Acknowledged by
Principal



Muslim Likabu, S.Pd
NIP. 19690718 200012 1 002

Timbong, 03 December 2021

English Teacher

Enrimon Piisi, S.Pd
NIP. 19870903 201101 1 007

Student Reflection



let's reflect!

Put a tick (✓) in the right column to show your feeling.
Berikan tanda centang (✓) pada kolom yang tepat!



= YEAH



= NO

No.	Question		
1.	Do you like to learn about Medicine Label?	✓	
2.	Are the pictures interesting?	✓	
3.	Is it easy to understand the lesson?	✓	
4.	Are there too many new words?	✓	
5.	Do you want to learn more about it?	✓	

Teacher Reflection

Medicine Label

1. What did I change from the original materials? Why?

At the next meeting I will adjust the position of the drug label, so that it is easier for students to read.

2. Are the materials you have developed appropriate to your students? Or are they only appropriate to a certain group of students in your class?

In my opinion, these food and beverage label materials are appropriate, but they still need to be further developed to make it more optimal.

3. What do you like most about the materials you develop? Why?

I really like this material when students will choose the right drug label for sick people, because I think this is an important thing.

4. From 1-10, how do you rate your materials? Why?

8 because, this material is the main material, which I have tried to make it easier for students to work on their LKPD. Improvements must still be made regarding LCD PROJECTORS that do not yet have complementary screens, in the learning process they still use blackboards and walls as replacement tools for displaying PPT.