

RENCANA PELAKSANAAN PEMBELAJARAN

SATUAN PEMBELAJARAN	: SMP NEGERI 1 Sangasanga
Mata Pelajaran	: Bahasa Inggris
Kelas / Semester	: IX / Ganjil
Materi Pokok	: Drug Label
Alokasi Waktu	: 10 Menit

A. Kompetensi Dasar

- 3.3 Membandingkan fungsi sosial, struktur teks, dan unsur kebahasaan beberapa teks khusus dalam bentuk label, dengan meminta dan memberi informasi terkait obat/makanan/minuman, sesuai dengan konteks penggunaannya
- 4.3 menangkap makna secara kontekstual terkait dengan fungsi sosial, struktur teks, dan unsur kebahasaan teks khusus dalam bentuk label pendek dan sederhana, terkait obat/makanan/minuman

B. Tujuan Pembelajaran

- Peserta didik dapat mengidentifikasi fungsi sosial, struktur teks, dan unsur kebahasaan dari teks *label* dengan meminta dan memberi informasi terkait obat sesuai dengan konteks penggunaannya
- Peserta didik dapat menemukan informasi tersirat dan tersurat dari teks *Drug Label*.

C. Langkah-langkah Kegiatan Pembelajaran

Alat : Gambar Drug's label & LKPD

Bahan : Internet (<https://www.drugwatch.com/health/how-to-read-a-drug-label/>)

1. Kegiatan Pembuka

- Guru membuka pembelajaran dengan salam
- Guru menanyakan kabar peserta didik dan memastikan kehadiran peserta didik
- Guru memimpin doa sebelum memulai pembelajaran
- Guru menanyakan kembali tentang pembelajaran sebelumnya (*Food Label*)
- Guru menyampaikan tujuan pembelajaran pada pertemuan tersebut

2. Kegiatan Inti

- Peserta didik bekerja berpasangan untuk memasangkan kartu frasa dan kartu definisi (*Pre-Teach Vocabulary*)
- Peserta didik membaca teks dan meletakkan frase / *Drug Facts Phrase* kedalam teks rumpang dengan benar. (LKPD 1)
- Peserta didik menjawab beberapa pertanyaan dari guru (lisan) terkait fungsi sosial drug label. (Guru memancing/*eliciting* jawaban dari peserta didik)
- Peserta didik membaca dan menjawab pertanyaan-pertanyaan pada LKPD 2 secara individu.
- Peserta didik mengecek jawabannya dengan teman disampingnya (*Peer Check*).
- Guru mengecek jawaban peserta didik secara bersama-sama dan memberikan umpan balik (*whole class feedback, nominate, pounce & bounce*)

3. Kegiatan Penutup

- Peserta didik menyimpulkan pembelajaran terkait dengan drug label
- Peserta didik mengungkapkan perasaannya setelah mengikuti pembelajaran hari ini
- Guru menyampaikan rencana pembelajaran berikutnya
- Guru menutup pembelajaran dengan salam

D. Penilaian Pembelajaran

- Guru menilai LKPD
- Guru merekap penilaian sikap peserta didik selama mengikuti pembelajaran

Mengetahui,
Kepala Sekolah

Sangasanga, 6 Januari 2022

Guru Mata Pelajaran

Rebo, S.Pd
NIP 196406151988031016

Rezha Hardi Pratama, S.Pd
NIP. -

Lampiran

Pre-teaching Vocabulary

Work in pair. Match the phrase and the definition correctly. In 5 minutes! (cut-phrases)

Phrase	Definition
Active Ingredient	Bahan aktif penyusun obat yang membuat obat bekerja efektif
Uses	Gejala-gejala penyakit yang diobati
Warnings	Peringatan, Efek samping
Directions	Jumlah (dosis) obat, seberapa sering dikonsumsi, dan seberapa banyak obat tersebut dapat dikonsumsi secara aman dalam sehari.
Other Information	Hal-hal penting untuk diketahui seperti bagaimana cara penyimpanan obat tersebut.
Inactive Ingredients	Rasa obat, bahan pengawet, atau bahan lain yang tidak berfungsi untuk mengobati gejala penyakit.
Questions / Comments	Nomor telepon perusahaan yang dapat dihubungi jika memiliki pertanyaan / komentar tentang obat tersebut.

LKPD 1

In pairs, read the drug label and fill in the gaps with suitable phrase given.

1	2	3	4	5	6	7								
1	2	3	4	5	6	7								
Purpose (in each caplet) Acetaminophen 500 mg.....Pain reliever/fever reducer	Drug Facts (continued) Stop use and ask a doctor if <ul style="list-style-type: none"> ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present These could be signs of a serious condition.	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	■ do not take more than directed (see overdose warning) <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="padding: 5px;">adults and children 12 years and over</td> <td style="padding: 5px;">■ take 2 caplets every 6 hours while symptoms last</td> </tr> <tr> <td style="padding: 5px;"></td> <td style="padding: 5px;">■ do not take more than 6 caplets in 24 hours, unless directed by a doctor</td> </tr> <tr> <td style="padding: 5px;"></td> <td style="padding: 5px;">■ do not use for more than 10 days unless directed by a doctor</td> </tr> </table> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="padding: 5px;">children under 12 years</td> <td style="padding: 5px;">ask a doctor</td> </tr> </table>	adults and children 12 years and over	■ take 2 caplets every 6 hours while symptoms last		■ do not take more than 6 caplets in 24 hours, unless directed by a doctor		■ do not use for more than 10 days unless directed by a doctor	children under 12 years	ask a doctor	■ store at 20-25°C (68-77°F)	carnauba wax, corn starch*, croscarmellose sodium , hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid *may contain one or more of these ingredients	1-800-719-9260
adults and children 12 years and over	■ take 2 caplets every 6 hours while symptoms last													
	■ do not take more than 6 caplets in 24 hours, unless directed by a doctor													
	■ do not use for more than 10 days unless directed by a doctor													
children under 12 years	ask a doctor													
Active Ingredients	Inactive ingredients	Other Information	Warnings	Uses	Questions / Comments	Directions								

Source: <https://fda.report/DailyMed/c9d3cee1-531c-439f-ae19-07acd366aad6>

Act

Active Ingredients	Inactive ingredients	Other Information	Warnings	Uses	Questions / Comments	Directions

Individually, read the label and the statements carefully.
Write **T** if you think the statement is true and **F** if it is false.

1	Drug Facts	Drug Facts (continued)				
Active ingredient (in each caplet)	Purpose	Stop use and ask a doctor if				
Acetaminophen 500 mg.....	Pain reliever/fever reducer	<ul style="list-style-type: none"> ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present <p>These could be signs of a serious condition.</p>				
2	Uses	If pregnant or breast-feeding, ask a health professional before use.				
<ul style="list-style-type: none"> ■ temporarily relieves minor aches and pains due to: ■ the common cold ■ headache ■ minor pain of arthritis ■ backache ■ muscular aches ■ toothache ■ premenstrual and menstrual cramps ■ temporarily reduces fever 	Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	3				
Warnings	Directions	4				
<p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> ■ skin reddening ■ blisters ■ rash <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</p> <ul style="list-style-type: none"> ■ if you have ever had an allergic reaction to this product or any of its ingredients <p>Ask a doctor before use if you have liver disease</p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin</p>	<p>■ do not take more than directed (see overdose warning)</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%;">adults and children 12 years and over</td> <td style="width: 50%;"> <ul style="list-style-type: none"> ■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor </td> </tr> <tr> <td>children under 12 years</td> <td>ask a doctor</td> </tr> </table>	adults and children 12 years and over	<ul style="list-style-type: none"> ■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor 	children under 12 years	ask a doctor	5
adults and children 12 years and over	<ul style="list-style-type: none"> ■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor 					
children under 12 years	ask a doctor					
	Other information	6				
	<p>Inactive ingredients carnauba wax, corn starch*, croscarmellose sodium*, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid. *may contain one or more of these ingredients</p>	7				
	Questions or comments? 1-800-719-9260					

Act

Source: <https://fda.report/DailyMed/c9d3cee1-531c-439f-ae19-07acd366aad6>

No.	Statement	T / F
1	The product contains more than 4000 mg acetaminophen.	
2	Acetaminophen may cause severe skin reactions.	
3	We must store the product between 68 – 77 degree of Celcius	
4	12-year-old child must ask a doctor before consuming the product.	
5	We must stop using the product if the pain and fever get worse.	
6	We can consume it with any other drug containing acetaminophen	
7	Consuming warfarin with this medicine may be dangerous to some people	
8	Adults can consume the medicine without ask a doctor first	
9	Sodium starch glycolate in it have a medical purpose	
10	Call 1-900-719-9250 if overdose is happened	